

Stability of SARS-CoV-2 in different environmental conditions

Alex W.H. Chin, Julie T.S. Chu, Mahen R.A. Perera, Kenrie P.Y. Hui, Hui-Ling Yen, Michael C.W. Chan, Malik Peiris, Leo L.M. Poon*

School of Public Health, LKS Faculty of Medicine, The University of Hong Kong, Hong Kong Special Administrative Region, China.

* Corresponding authors:

- Leo Poon, School of Public Health, LKS Faculty of Medicine, The University of Hong Kong, Hong Kong SAR, China. E-mail: llmpoon@hku.hk

To the Editor,

We previously reported the detection of SARS-CoV-2 in different clinical samples¹. This virus can be detected on different surfaces in a contaminated site². Here, we report the stability of SARS-CoV-2 in different environmental conditions.

We first determined the stability of SARS-CoV-2 at different temperatures. SARS-CoV-2 in virus transport medium (VTM; final concentration: ~ 6.8 log TCID₅₀/mL) was incubated for up to 14 days and then tested for its infectivity (Table A). The virus is highly stable at 4°C, but sensitive to heat. At 4°C, there was only ~ 0.7 -log unit reduction of infectious titre on Day 14. With the incubation temperature being increased to 70°C, the time for virus inactivation was reduced to 5 minutes.

We further investigated the stability of this virus on different surfaces. In brief, a 5- μ L droplet of virus culture (~ 7.8 Log unit of TCID₅₀/mL) was pipetted on a surface (Table B; $\sim 1\text{cm}^2$ per piece) and left at room temperature (22°C; Relative humidity: $\sim 65\%$). The inoculated objects retrieved at desired time points were immediately soaked with 200 μ L of VTM for 30 minutes to elute the virus. No infectious virus could be recovered from printing and tissue papers after a 3-hour incubation, whereas no infectious virus could be detected from treated wood and cloth on Day 2. By contrast, SARS-CoV-2 was more stable on smooth surfaces. No infectious virus could be detected from treated smooth surfaces on Day 4 (glass and banknote) or Day 7 (stainless steel and plastic). Strikingly, a significant level of infectious virus could still be detected on the outer layer of a surgical mask on Day 7 ($\sim 0.1\%$ of the original inoculum). Interestingly, a biphasic decay of infectious SARS-CoV-2 could be found from samples recovered from these smooth surfaces (Appendix). Representative negative samples were tested positive by RT-PCR³ (N=39; data not shown), demonstrating that non-infectious viruses could be recovered by the eluents.

We also tested the virucidal effects of disinfectants by adding 15 μ L of SARS-CoV-2 culture (~ 7.8 Log unit of TCID₅₀/mL) to 135 μ L of various disinfectants at working concentration (Table C). With the exception of a 5-min incubation with hand soap, no infectious virus could be detected after a 5-minute incubation at room temperature. In addition, we also found that

SARS-CoV-2 is extremely stable in a wide-range of pH values at room temperature (pH3-10; Table D)

Overall, SARS-CoV-2 can be highly stable in a favourable environment⁴, but it is also susceptible to standard disinfection methods.

We declare that we have no competing interests. This work was supported by NIADI, NIH (USA) (contract HHSN272201400006C). LLMP was supported by Croucher Foundation.

References:

1. Pan Y, Zhang D, Yang P, Poon LLM, Wang Q. Viral load of SARS-CoV-2 in clinical samples. Lancet Infect Dis (2020) published online Feb 24. [https://www.thelancet.com/journals/laninf/article/PIIS1473-3099\(20\)30113-4/fulltext](https://www.thelancet.com/journals/laninf/article/PIIS1473-3099(20)30113-4/fulltext)
2. Ye G, Lin H, Chen L. et al. Environmental contamination of the SARS-CoV-2 in healthcare premises: An urgent call for protection for healthcare workers. medRxiv 2020.03.11.20034546; doi: <https://doi.org/10.1101/2020.03.11.20034546>
3. Chu DKW, Pan Y, Cheng SMS, et al. Molecular Diagnosis of a Novel Coronavirus (2019-nCoV) Causing an Outbreak of Pneumonia. Clin Chem (2020) published online Jan 31. <https://academic.oup.com/clinchem/advance-article/doi/10.1093/clinchem/hvaa029/5719336>
4. van Doremalen N, Bushmaker T, Morris DH, et al. Aerosol and Surface Stability of SARS-CoV-2 as Compared with SARS-CoV-1. N Engl J Med (2020) published online March 17. https://www.nejm.org/doi/full/10.1056/NEJMc2004973?query=featured_home

Table. Stability of SARS-CoV-2 at different environmental conditions.

Time	Virus titre (Log TCID ₅₀ /mL)									
	4°C		22°C		37°C		56°C		70°C	
	Mean	±SD	Mean	±SD	Mean	±SD	Mean	±SD	Mean	±SD
1 min	N.D.	N.D.	6.51	0.27	N.D.	N.D.	6.65	0.1	5.34	0.17
5 mins	N.D.	N.D.	6.7	0.15	N.D.	N.D.	4.62	0.44	U	-
10 mins	N.D.	N.D.	6.63	0.07	N.D.	N.D.	3.84	0.32	U	-
30 mins	6.51	0.27	6.52	0.28	6.57	0.17	U	-	U	-
1 hr	6.57	0.32	6.33	0.21	6.76	0.05	U	-	U	-
3 hrs	6.66	0.16	6.68	0.46	6.36	0.19	U	-	U	-
6 hrs	6.67	0.04	6.54	0.32	5.99	0.26	U	-	U	-
12 hrs	6.58	0.21	6.23	0.05	5.28	0.23	U	-	U	-
1 day	6.72	0.13	6.26	0.05	3.23	0.05	U	-	U	-
2 days	6.42	0.37	5.83	0.28	U	-	U	-	U	-
4 days	6.32	0.27	4.99	0.18	U	-	U	-	U	-
7 days	6.65	0.05	3.48	0.24	U	-	U	-	U	-
14 days	6.04	0.18	U	-	U	-	U	-	U	-

A) Temperature*

B) Surfaces*

Time	Virus titre (Log TCID ₅₀ /mL)									
	Paper		Tissue paper		Wood		Cloth		Glass	
	Mean	±SD	Mean	±SD	Mean	±SD	Mean	±SD	Mean	±SD
0 min	4.76	0.10	5.48	0.10	5.66	0.39	4.84	0.17	5.83	0.04
30 mins	2.18	0.05	2.19	0.17	3.84	0.39	2.84	0.24	5.81	0.27
3 hrs	U	-	U	-	3.41	0.26	2.21"	-	5.14	0.05
6 hrs	U	-	U	-	2.47	0.23	2.25	0.08	5.06	0.31
1 day	U	-	U	-	2.07"	-	2.07"	-	3.48	0.37
2 days	U	-	U	-	U	-	U	-	2.44	0.19
4 days	U	-	U	-	U	-	U	-	U	-
7 days	U	-	U	-	U	-	U	-	U	-

Time	Banknote		Stainless steel		Plastic		Mask, inner layer		Mask, outer layer	
	Mean	±SD	Mean	±SD	Mean	±SD	Mean	±SD	Mean	±SD
0 min	6.05	0.34	5.80	0.02	5.81	0.03	5.88	0.69	5.78	0.10
30 mins	5.83	0.29	5.23	0.05	5.83	0.04	5.84	0.18	5.75	0.08
3 hrs	4.77	0.07	5.09	0.04	5.33	0.22	5.24	0.08	5.11	0.29
6 hrs	4.04	0.29	5.24	0.08	4.68	0.10	5.01	0.50	4.97	0.51
1 day	3.29	0.60	4.85	0.20	3.89	0.33	4.21	0.08	4.73	0.05
2 days	2.47	0.23	4.44	0.20	2.76	0.10	3.16	0.07	4.20	0.07
4 days	U	-	3.26	0.10	2.27	0.09	2.47	0.28	3.71	0.50
7 days	U	-	U	-	U	-	U	-	2.79	0.46

C) Disinfectants*

Disinfectant (Working concentration)	Virus titre (Log TCID ₅₀ /mL)		
	5 mins	15 mins	30 mins
Household bleach (1:49)	U	U	U
Household bleach (1:99)	U	U	U
Hand soap solution (1:49)	3.6"	U	U
Ethanol (70%)	U	U	U
Povidone-iodine (7.5%)	U	U	U
Chloroxylonol (0.05%)	U	U	U
Chlorhexidine (0.05%)	U	U	U
Benzalkonium chloride (0.1%)	U	U	U

D) pH*

pH (60 mins)	Virus titre (Log TCID ₅₀ /mL)	
	Mean	±SD
3	5.55	0.25
4	5.67	0.36
5	5.73	0.04
6	5.75	0.08
7	5.58	0.22
8	5.70	0.14
9	5.54	0.44
10	5.51	0.11

* All the virus titres were titrated using Vero-E6 cells. All experimental studies were done in three independent triplicates. Detection limit of a typical TCID₅₀ assay is 100 TCID₅₀/mL, except reactions containing hand soap/chloroxylonol (detection

limit: 10^3 TCID₅₀/mL) or reactions containing povidone-iodine/chlorhexidin/benzalkonium chloride; detection limit: 10^4 TCID₅₀/mL) because of their cytotoxic effects. N.D.: not done, U: undetectable.
Only one of the triplicate reactions was positive in the TCID₅₀ assay.