

Cite as: K. W. Ng *et al.*, *Science*  
10.1126/science.abe1107 (2020).

# Preexisting and de novo humoral immunity to SARS-CoV-2 in humans

Kevin W. Ng<sup>1\*</sup>, Nikhil Faulkner<sup>1\*</sup>, Georgina H. Cornish<sup>1\*</sup>, Annachiara Rosa<sup>2\*</sup>, Ruth Harvey<sup>3</sup>, Saira Hussain<sup>3</sup>, Rachel Ulferts<sup>9</sup>, Christopher Earl<sup>4</sup>, Antoni G. Wrobel<sup>5</sup>, Donald J. Benton<sup>5</sup>, Chloe Roustan<sup>6</sup>, William Bolland<sup>1</sup>, Rachael Thompson<sup>1</sup>, Ana Agua-Doce<sup>7</sup>, Philip Hobson<sup>7</sup>, Judith Heaney<sup>13</sup>, Hannah Rickman<sup>13</sup>, Stavroula Paraskevopoulou<sup>13</sup>, Catherine F. Houlihan<sup>13,14</sup>, Kirsty Thomson<sup>13</sup>, Emilie Sanchez<sup>13</sup>, Gee Yen Shin<sup>13</sup>, Moira J. Spyer<sup>13,15</sup>, Dhira Joshi<sup>8</sup>, Nicola O'Reilly<sup>8</sup>, Philip A. Walker<sup>6</sup>, Svend Kjaer<sup>6</sup>, Andrew Riddell<sup>7</sup>, Catherine Moore<sup>16</sup>, Bethany R. Jebson<sup>17,19</sup>, Meredyth Wilkinson<sup>17,19</sup>, Lucy R. Marshall<sup>17,19</sup>, Elizabeth C. Rosser<sup>17,18</sup>, Anna Radziszewska<sup>17,18</sup>, Hannah Peckham<sup>17,18</sup>, Coziana Ciurtin<sup>17,18</sup>, Lucy R. Wedderburn<sup>17,19</sup>, Rupert Beale<sup>9</sup>, Charles Swanton<sup>10</sup>, Sonia Gandhi<sup>11</sup>, Brigitta Stockinger<sup>12</sup>, John McCauley<sup>3</sup>, Steve J. Gamblin<sup>5</sup>, Laura E. McCoy<sup>14†</sup>, Peter Cherepanov<sup>2†</sup>, Eleni Nastouli<sup>13,15†</sup>, George Kassiotis<sup>1,20†</sup>

<sup>1</sup>Retroviral Immunology, The Francis Crick Institute, London NW1 1AT, UK. <sup>2</sup>Chromatin Structure and Mobile DNA Laboratory, The Francis Crick Institute, London NW1 1AT, UK. <sup>3</sup>Worldwide Influenza Centre, The Francis Crick Institute, London NW1 1AT, UK. <sup>4</sup>Signalling and Structural Biology Laboratory, The Francis Crick Institute, London NW1 1AT, UK. <sup>5</sup>Structural Biology of Disease Processes Laboratory, The Francis Crick Institute, London NW1 1AT, UK. <sup>6</sup>Structural Biology STP, The Francis Crick Institute, London NW1 1AT, UK. <sup>7</sup>Flow Cytometry STP, The Francis Crick Institute, London NW1 1AT, UK. <sup>8</sup>Peptide Chemistry, The Francis Crick Institute, London NW1 1AT, UK. <sup>9</sup>Cell Biology of Infection Laboratory, The Francis Crick Institute, London NW1 1AT, UK. <sup>10</sup>Cancer Evolution and Genome Instability Laboratory, The Francis Crick Institute, London NW1 1AT, UK. <sup>11</sup>Neurodegeneration Biology Laboratory, The Francis Crick Institute, London NW1 1AT, UK. <sup>12</sup>AhRimmunity Laboratory, The Francis Crick Institute, London NW1 1AT, UK. <sup>13</sup>University College London Hospitals (UCLH) NHS Trust, London NW1 2BU, UK. <sup>14</sup>Division of Infection and Immunity, University College London (UCL), London WC1E 6BT, UK. <sup>15</sup>Department of Population, Policy and Practice, Great Ormond Street Institute for Child Health (ICH), UCL, London WC1N 1EH, UK. <sup>16</sup>Public Health Wales, University Hospital of Wales, Cardiff CF14 4XW, UK. <sup>17</sup>Centre for Adolescent Rheumatology Versus Arthritis at UCL, UCLH, Great Ormond Street Hospital (GOSH), London WC1N 3JH, UK. <sup>18</sup>Centre for Rheumatology Research, Division of Medicine, UCL, London, WC1E 6BT, UK. <sup>19</sup>UCL Great Ormond Street Institute for Child Health (ICH), UCL, London WC1N 1EH, UK. <sup>20</sup>Department of Medicine, Faculty of Medicine, Imperial College London, London W2 1PG, UK.

\*These authors contributed equally to this work.

†Corresponding author. Email: george.kassiotis@crick.ac.uk (G.K.); e.nastouli@ucl.ac.uk (E.N.); peter.cherepanov@crick.ac.uk (P.C.); l.mccoy@ucl.ac.uk (L.E.M.)

**Zoonotic introduction of novel coronaviruses may encounter preexisting immunity in humans. Using diverse assays for antibodies recognizing SARS-CoV-2 proteins, we detect preexisting humoral immunity. SARS-CoV-2 spike glycoprotein (S)-reactive antibodies were detectable by a flow cytometry-based method in SARS-CoV-2-uninfected individuals and were particularly prevalent in children and adolescents. They were predominantly of the IgG class and targeted the S2 subunit. By contrast, SARS-CoV-2 infection induced higher titers of SARS-CoV-2 S-reactive IgG antibodies, targeting both the S1 and S2 subunits, and concomitant IgM and IgA antibodies, lasting throughout the observation period. Notably, SARS-CoV-2-uninfected donor sera exhibited specific neutralizing activity against SARS-CoV-2 and SARS-CoV-2 S pseudotypes. Distinguishing preexisting and de novo immunity will be critical for our understanding of susceptibility to and the natural course of SARS-CoV-2 infection.**

Immune cross-reactivity among seasonally spreading human coronaviruses (HCoV) has long been hypothesized to provide effective, but transient cross-protection against distinct HCoVs (1, 2). To determine the degree of cross-reactivity between HCoVs and SARS-CoV-2, we developed a flow cytometry-based assay for SARS-CoV-2-binding antibodies. The main target for such antibodies is the spike glycoprotein (S), which is proteolytically processed into the S1 and S2 subunits, mediating target cell attachment and entry, respectively.

The S1-specific CR3022 antibody stained a smaller percentage of SARS-CoV-2 S-expressing HEK293T cells and with lower intensity than COVID-19 convalescent sera (fig. S1), indicating that polyclonal IgG antibodies targeted a wider range of epitopes naturally processed and displayed on these

cells. This assay also detected SARS-CoV-2 S-reactive IgM and IgA antibodies in COVID-19 convalescent sera (fig. S2). Indeed, the presence of SARS-CoV-2 S-reactive antibodies of all three Ig classes (IgG<sup>+</sup>IgM<sup>+</sup>IgA<sup>+</sup>) distinguished COVID-19 sera from control sera with a high degree of sensitivity and specificity (Fig. 1A and fig. S3). All 156 seroconverted COVID-19 patients had contemporaneous IgG, IgM, and IgA responses to SARS-CoV-2 S throughout the observation period, with the exception of two patients, who only had IgG antibodies (figs. S4 and S5). One of these patients was a bone-marrow-transplant recipient who experienced HCoV infection a month prior to SARS-CoV-2 infection (fig. S6). Remarkably, a small proportion of SARS-CoV-2-uninfected patients, sampled before or during the early spread of SARS-CoV-2 in the UK

(table S1), also had SARS-CoV-2 S-binding IgG, but not IgM or IgA antibodies (Fig. 1A), suggesting the presence of cross-reactive immunological memory.

The S2 subunit exhibits a higher degree of homology among coronaviruses than S1 (fig. S7) and was likely the main target of cross-reactive antibodies. Competition with recombinant soluble S1 or S2, at doses that blocked binding of specific monoclonal antibodies (fig. S8), did not affect the frequency of cells stained with COVID-19 patient sera, although the intensity of staining was reduced by 31% and 37%, respectively (Fig. 1, B to D), indicating recognition of both S1 and S2. By contrast, soluble S2 completely abolished staining with SARS-CoV-2-uninfected patient sera, whereas soluble S1 had no effect (Fig. 1, B to D). Thus, SARS-CoV-2-uninfected patient sera cross-reacts with SARS-CoV-2 S2 and COVID-19 patient sera additionally recognizes S1.

SARS-CoV-2 S-reactive IgG antibodies were detected by flow cytometry in 5 of 34 SARS-CoV-2-uninfected individuals with RT-qPCR-confirmed HCoV infection, as well as in 1 of 31 individuals without recent HCoV infection (Fig. 2A and fig. S4A). This suggested that cross-reactivity may have persisted from earlier HCoV infections, rather than having been induced by the most recent one.

To confirm antibody cross-reactivity using an independent assay, we developed ELISAs using recombinant SARS-CoV-2 stabilized trimeric S ectodomain, S1, receptor-binding domain (RBD), or nucleoprotein (N). Rates of IgG seropositivity by SARS-CoV-2 S1-coated ELISA were congruent with, but generally lower than those by flow cytometry (fig. S9). The three SARS-CoV-2-uninfected individuals with the highest cross-recognition of S by flow cytometry, plus an additional four individuals, had ELISA-detectable IgG antibodies against SARS-CoV-2 S ectodomain, as well as N (Fig. 2A and fig. S4, B to D). By contrast, none of the control samples had ELISA-detectable IgG antibodies against the less conserved SARS-CoV-2 S1 or RBD (Fig. 2A and fig. S4, B to D).

The prevalence of such cross-reactive antibodies was further examined in additional healthy donor cohorts (table S1). Among 50 SARS-CoV-2-uninfected pregnant women, sampled in May 2018, five showed evidence for SARS-CoV-2 S-reactive IgG, but not IgM or IgA antibodies (Fig. 2B and fig. S10). In a separate cohort of 101 SARS-CoV-2-uninfected donors, sampled in May 2019, three had SARS-CoV-2 S-reactive IgG antibodies (fig. S11), which did not correlate with antibodies to diverse viruses and bacteria also present in several of these samples. SARS-CoV-2 S-reactive IgM and IgA were also detected in two of these donors, albeit at considerably lower levels than in COVID-19 patients (fig. S11), suggestive of recent or ongoing response. In an additional cohort of 13 donors recently infected with HCoVs, only one had SARS-CoV-2 S-reactive IgG antibodies, and these at very low levels (fig. S12). This suggested that their emergence was not simply

a common transient event following each HCoV infection in this age group (median age 51 years; table S1). Instead, given that HCoV-reactive antibodies are present in virtually all adults (3–5), the rarity of SARS-CoV-2 S cross-reactivity (16 of 302; 5.29%), indicates additional requirements, such as random B cell receptor repertoire focusing or frequency of HCoV infection, rather than time since the last HCoV infection. Indeed, the frequency of HCoV infection displays a characteristic age distribution, being the highest in children and adolescents (1, 4–8). We therefore examined a cohort of younger SARS-CoV-2-uninfected healthy donors (aged 1–16 years; table S1), sampled between 2011 and 2018. Remarkably, at least 21 of these 48 donors had detectable levels of SARS-CoV-2 S-reactive IgG antibodies (Fig. 2, C to E), whereas only one of an additional cohort of 43 young adults (aged 17–25 years; table S1) had such antibodies (Fig. 2F). Staining with sera from SARS-CoV-2-uninfected children and adolescents was specific to HEK293T cells expressing SARS-CoV-2 S, but not the unrelated HERV-K113 envelope glycoprotein, and was outcompeted by soluble SARS-CoV-2 S2 (fig. S13). The prevalence of SARS-CoV-2 S-reactive IgG antibodies peaked at 62% between 6 and 16 years of age (Fig. 2F), when HCoV seroconversion in this age group also peaks (3, 4, 6, 7), and was significantly higher than in adults ( $P < 0.00001$ , Fisher's exact test).

To probe the potential consequences of antibody cross-reactivity, we examined the ability of preexisting antibodies to inhibit SARS-CoV-2 entry into HEK293T cells (fig. S14 and supplementary text). Although not expected to directly inhibit RBD-mediated cell attachment, S2-targeting antibodies that can neutralize SARS-CoV-2 have recently been discovered (9, 10). HEK293T cell infection with SARS-CoV-2 S pseudotypes was efficiently inhibited by sera from seroconverted ( $Ab^+$ ) COVID-19 patients, but not from those that had not yet seroconverted ( $Ab^-$ ) (Fig. 3A). Surprisingly, sera from SARS-CoV-2-uninfected donors with SARS-CoV-2 S-reactive antibodies also neutralized these pseudotypes, whereas none of the sera neutralized vesicular stomatitis virus (VSV) glycoprotein pseudotypes (Fig. 3A). Comparable neutralization of SARS-CoV-2 S pseudotypes was also observed with sera from SARS-CoV-2-uninfected adolescents (Fig. 3A). Moreover, the majority of sera from SARS-CoV-2-uninfected donors with flow cytometry-detectable cross-reactive antibodies also neutralized authentic SARS-CoV-2 infection of Vero E6 cells, albeit on average less potently than COVID-19 patient sera (Fig. 3B). By contrast, sera from SARS-CoV-2-uninfected patients without cross-reactive antibodies exhibited no neutralizing activity (Fig. 3B). Antiviral antibodies may also enhance viral entry by Fc receptor-mediated antibody-dependent enhancement (ADE). However, entry of SARS-CoV-2 S pseudotypes was not enhanced by either COVID-19 patient or SARS-CoV-2-uninfected patient sera in Fc $\gamma$ RIIA-expressing K-562 cells (fig. S15).

Collectively, these findings highlighted functionally relevant antigenic epitopes conserved within the S2. Over its entire length, SARS-CoV-2 S exhibits marginally closer homology with the S proteins of betacoronaviruses HCoV-OC43 and HCoV-HKU1, than of alphacoronaviruses HCoV-NL63 and HCoV-229E (fig. S16A). To probe shared epitopes, we constructed overlapping peptide arrays spanning the last 743 amino acids of SARS-CoV-2 S (fig. S16B). Multiple putative epitopes were differentially recognized by sera with cross-reactive antibodies (Ab<sup>+</sup>), were reasonably conserved, and most mapped to the surface of S2 (Fig. 4, A and B, and table S2). Of note, an epitope overlapping the S2 fusion peptide was also recently identified as cross-reactive with the corresponding peptides from HCoV-OC43 and HCoV-229E (11). Cross-reactivity with the identified epitopes was further supported by ELISAs coated with synthetic peptides (fig. S17).

As expected (3–5), reactivity with one or more HCoVs was detectable by flow cytometry in all sera (Fig. 4D and fig. S18). However, IgG and IgA reactivity against HCoVs was higher in SARS-CoV-2-uninfected adults with, than without SARS-CoV-2-reactive IgG ( $P=1.4\times 10^{-6}$  for IgG and  $P=0.017$  for IgA, Student's *t* test), and in SARS-CoV-2-uninfected children or adolescents with, than without SARS-CoV-2-reactive IgG ( $P=0.010$  for IgG and  $P=0.021$  for IgA, Student's *t* test) (Fig. 4D), supporting a direct link between the two. Accordingly, IgG reactivity against each HCoV type was independently correlated with the presence of SARS-CoV-2-reactive antibodies (Fig. 4D).

Our results from multiple independent assays demonstrated the presence of preexisting antibodies recognizing SARS-CoV-2 in uninfected individuals. Identification of conserved epitopes in S2 targeted by neutralizing antibodies may hold promise for a universal vaccine protecting against current, as well as future CoVs. Together with preexisting T cell (12–14) and B cell memory (10, 15), antibody cross-reactivity between seasonal HCoVs and SARS-CoV-2 may have important ramifications for natural infection. Epidemiological studies of HCoV transmission suggest that cross-protective immunity is unlikely to be sterilizing or long-lasting (8), which is also supported by repeated reinfection (2, 16). Nevertheless, prior immunity induced by one HCoV can reduce the transmission of homologous and heterologous HCoVs, and ameliorate the symptoms where transmission is not prevented (1, 2). A possible modification of COVID-19 severity by prior HCoV infection may account for the age distribution of COVID-19 susceptibility, where higher HCoV infection rates in children than in adults (4, 6) correlates with relative protection from COVID-19 (17) and may also shape seasonal and geographical patterns of transmission. It is, therefore, imperative that any effect, positive or negative, of preexisting HCoV-elicited immunity on the natural course of SARS-CoV-2 infection be fully delineated.

## REFERENCES AND NOTES

1. R. W. Aldridge, D. Lewer, S. Beale, A. M. Johnson, M. Zambon, A. C. Hayward, E. B. Fragaszy, Seasonality and immunity to laboratory-confirmed seasonal coronaviruses (HCoV-NL63, HCoV-OC43, and HCoV-229E): Results from the Flu

Watch cohort study. *Wellcome Open Res.* 5, 52 (2020).

[doi:10.12688/wellcomeopenres.15812.1](https://doi.org/10.12688/wellcomeopenres.15812.1)

2. K. A. Callow, H. F. Parry, M. Sergeant, D. A. Tyrrell, The time course of the immune response to experimental coronavirus infection of man. *Epidemiol. Infect.* 105, 435–446 (1990). [doi:10.1017/S0950268800048019](https://doi.org/10.1017/S0950268800048019) [Medline](#)
3. E. G. Severance, I. Bossis, F. B. Dickerson, C. R. Stallings, A. E. Origoni, A. Sullens, R. H. Yolken, R. P. Viscidi, Development of a nucleocapsid-based human coronavirus immunoassay and estimates of individuals exposed to coronavirus in a U.S. metropolitan population. *Clin. Vaccine Immunol.* 15, 1805–1810 (2008). [doi:10.1128/CVI.00124-08](https://doi.org/10.1128/CVI.00124-08) [Medline](#)
4. R. Dijkman, M. F. Jebbink, N. B. El Idrissi, K. Pyrc, M. A. Müller, T. W. Kuijpers, H. L. Zaaijer, L. van der Hoek, Human coronavirus NL63 and 229E seroconversion in children. *J. Clin. Microbiol.* 46, 2368–2373 (2008). [doi:10.1128/JCM.00533-08](https://doi.org/10.1128/JCM.00533-08) [Medline](#)
5. A. T. Huang, B. Garcia-Carreras, M. D. T. Hitchings, B. Yang, L. C. Katzelnick, S. M. Rattigan, B. A. Borgert, C. A. Moreno, B. D. Solomon, L. Trimmer-Smith, V. Etienne, I. Rodriguez-Barraquer, J. Lessler, H. Salje, D. S. Burke, A. Wesolowski, D. A. T. Cummings, A systematic review of antibody mediated immunity to coronaviruses: Kinetics, correlates of protection, and association with severity. *Nat. Commun.* 11, 4704 (2020). [doi:10.1038/s41467-020-18450-4](https://doi.org/10.1038/s41467-020-18450-4) [Medline](#)
6. N. Friedman, H. Alter, M. Hindiyeh, E. Mendelson, Y. Shemer Avni, M. Mandelboim, Human coronavirus infections in Israel: Epidemiology, clinical symptoms and summer seasonality of HCoV-HKU1. *Viruses* 10, 515 (2018). [doi:10.3390/v10100515](https://doi.org/10.3390/v10100515) [Medline](#)
7. S. Nickbakhsh et al., Epidemiology of seasonal coronaviruses: Establishing the context for COVID-19 emergence. *J. Infect. Dis.* (2020).
8. A. S. Monto, P. M. DeJonghe, A. P. Callear, L. A. Bazzi, S. B. Capriola, R. E. Malosh, E. T. Martin, J. G. Petrie, Coronavirus occurrence and transmission over 8 years in the HIVE cohort of households in Michigan. *J. Infect. Dis.* 222, 9–16 (2020). [doi:10.1093/infdis/jiaa161](https://doi.org/10.1093/infdis/jiaa161) [Medline](#)
9. X. Chi, R. Yan, J. Zhang, G. Zhang, Y. Zhang, M. Hao, Z. Zhang, P. Fan, Y. Dong, Y. Yang, Z. Chen, Y. Guo, J. Zhang, Y. Li, X. Song, Y. Chen, L. Xia, L. Fu, L. Hou, J. Xu, C. Yu, J. Li, Q. Zhou, W. Chen, A neutralizing human antibody binds to the N-terminal domain of the Spike protein of SARS-CoV-2. *Science* 369, 650–655 (2020). [doi:10.1126/science.abc6952](https://doi.org/10.1126/science.abc6952) [Medline](#)
10. G. Song, W. He, S. Callaghan, F. Anzanello, D. Huang, J. Ricketts, J. L. Torres, N. Beutler, L. Peng, S. Vargas, J. Cassell, M. Parren, L. Yang, C. Ignacio, D. M. Smith, J. E. Voss, D. Nemazee, A. B. Ward, T. Rogers, D. R. Burton, R. Andrabi, Cross-reactive serum and memory B cell responses to spike protein in SARS-CoV-2 and endemic coronavirus infection. *bioRxiv* 308965 [Preprint]. 23 September 2020; <https://doi.org/10.1101/2020.09.22.308965>
11. E. Shrock, E. Fujimura, T. Kula, R. T. Timms, I.-H. Lee, Y. Leng, M. L. Robinson, B. M. Sie, M. Z. Li, Y. Chen, J. Logue, A. Zuiani, D. McCulloch, F. J. N. Lelis, S. Henson, D. R. Monaco, M. Travers, S. Habibi, W. A. Clarke, P. Caturegli, O. Laeyendecker, A. Piechocka-Trocha, J. Li, A. Khatir, H. Y. Chu, A.-C. Villani, K. Kays, M. B. Goldberg, N. Hacohen, M. R. Filbin, X. G. Yu, B. D. Walker, D. R. Wesemann, H. B. Larman, J. A. Lederer, S. J. Elledge; MGH COVID-19 Collection & Processing Team, Viral epitope profiling of COVID-19 patients reveals cross-reactivity and correlates of severity. *Science* eabd4250 (2020). [doi:10.1126/science.abd4250](https://doi.org/10.1126/science.abd4250) [Medline](#)
12. A. Grifoni, D. Weiskopf, S. I. Ramirez, J. Mateus, J. M. Dan, C. R. Moderbacher, S. A. Rawlings, A. Sutherland, L. Premkumar, R. S. Jodi, D. Marrama, A. M. de Silva, A. Frazier, A. F. Carlin, J. A. Greenbaum, B. Peters, F. Krammer, D. M. Smith, S. Crotty, A. Sette, Targets of T cell responses to SARS-CoV-2 coronavirus in humans with COVID-19 disease and unexposed individuals. *Cell* 181, 1489–1501.e15 (2020). [doi:10.1016/j.cell.2020.05.015](https://doi.org/10.1016/j.cell.2020.05.015) [Medline](#)
13. J. Braun, L. Loyal, M. Frentsch, D. Wendisch, P. Georg, F. Kurth, S. Hippenstiel, M. Dingeldey, B. Kruse, F. Fauchere, E. Baysal, M. Mangold, L. Henze, R. Lauster, M. A. Mall, K. Beyer, J. Röhm, S. Voigt, J. Schmitz, S. Miltenyi, I. Demuth, M. A. Müller, A. Hocke, M. Witzenth, N. Suttrop, F. Kern, U. Reimer, H. Wenschuh, C. Drosten, V. M. Corman, C. Giesecke-Thiel, L. E. Sander, A. Thiel, SARS-CoV-2-reactive T cells in healthy donors and patients with COVID-19. *Nature* (2020). [doi:10.1038/s41586-020-2598-9](https://doi.org/10.1038/s41586-020-2598-9) [Medline](#)
14. N. Le Bert, A. T. Tan, K. Kunasegaran, C. Y. L. Tham, M. Hafezi, A. Chia, M. H. Y. Chng, M. Lin, N. Tan, M. Linster, W. N. Chia, M. I.-C. Chen, L.-F. Wang, E. E. Ooi, S. Kalimuddin, P. A. Tambyah, J. G.-H. Low, Y.-J. Tan, A. Bertoletti, SARS-CoV-2



- specific T cell immunity in cases of COVID-19 and SARS, and uninfected controls. *Nature* **584**, 457–462 (2020). [doi:10.1038/s41586-020-2550-z](https://doi.org/10.1038/s41586-020-2550-z) [Medline](#)
15. P. Nguyen-Contant, A. K. Embong, P. Kanagaiah, F. A. Chaves, H. Yang, A. R. Branche, D. J. Topham, M. Y. Sangster, S protein-reactive IgG and memory B cell production after human SARS-CoV-2 infection includes broad reactivity to the S2 subunit. *mBio* **11**, e01991–e01920 (2020). [doi:10.1128/mBio.01991-20](https://doi.org/10.1128/mBio.01991-20) [Medline](#)
  16. P. K. Kiyuka, C. N. Agoti, P. K. Munywoki, R. Njeru, A. Bett, J. R. Otieno, G. P. Otieno, E. Kamau, T. G. Clark, L. van der Hoek, P. Kellam, D. J. Nokes, M. Cotten, Human coronavirus NL63 molecular epidemiology and evolutionary patterns in rural coastal Kenya. *J. Infect. Dis.* **217**, 1728–1739 (2018). [doi:10.1093/infdis/jiy098](https://doi.org/10.1093/infdis/jiy098) [Medline](#)
  17. R. Castagnoli, M. Votto, A. Licari, I. Brambilla, R. Bruno, S. Perlini, F. Rovida, F. Baldanti, G. L. Marseglia, Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection in children and adolescents: A systematic review. *JAMA Pediatr.* **174**, 882–889 (2020). [doi:10.1001/jamapediatrics.2020.1467](https://doi.org/10.1001/jamapediatrics.2020.1467) [Medline](#)
  18. P. R. Grant, M. A. Turner, G. Y. Shin, E. Nastouli, L. J. Levett, Extraction-free COVID-19 (SARS-CoV-2) diagnosis by RT-PCR to increase capacity for national testing programmes during a pandemic. *bioRxiv* 028316 [Preprint]. 9 April 2020; <https://doi.org/10.1101/2020.04.06.028316>.
  19. K. Hanke, P. Kramer, S. Seeher, N. Beiforde, R. Kurth, N. Bannert, Reconstitution of the ancestral glycoprotein of human endogenous retrovirus k and modulation of its functional activity by truncation of the cytoplasmic domain. *J. Virol.* **83**, 12790–12800 (2009). [doi:10.1128/JVI.01368-09](https://doi.org/10.1128/JVI.01368-09) [Medline](#)
  20. J. ter Meulen, E. N. van den Brink, L. L. M. Poon, W. E. Marissen, C. S. W. Leung, F. Cox, C. Y. Cheung, A. Q. Bakker, J. A. Bogaards, E. van Deventer, W. Preiser, H. W. Doerr, V. T. Chow, J. de Kruijff, J. S. M. Peiris, J. Goudsmit, Human monoclonal antibody combination against SARS coronavirus: Synergy and coverage of escape mutants. *PLOS Med.* **3**, e237 (2006). [doi:10.1371/journal.pmed.0030237](https://doi.org/10.1371/journal.pmed.0030237) [Medline](#)
  21. M. G. Joyce, R. S. Sankhala, W.-H. Chen, M. Choe, H. Bai, A. Hajducski, L. Yan, S. L. Sterling, C. E. Peterson, E. C. Green, C. Smith, N. de Val, M. Amare, P. Scott, E. D. Laing, C. C. Broder, M. Rolland, N. L. Michael, K. Modjarrad, A cryptic site of vulnerability on the receptor binding domain of the SARS-CoV-2 spike glycoprotein. *bioRxiv* 992883 [Preprint]. 17 March 2020; <https://doi.org/10.1101/2020.03.15.992883>.
  22. V. E. Pye, A. Rosa, C. Bertelli, W. B. Struwe, S. L. Maslen, R. Corey, I. Liko, M. Hassall, G. Mattiuzzo, A. Ballandras-Colas, A. Nans, Y. Takeuchi, P. J. Stansfeld, J. M. Skehel, C. V. Robinson, M. Pizzato, P. Cherepanov, A bipartite structural organization defines the SERINC family of HIV-1 restriction factors. *Nat. Struct. Mol. Biol.* **27**, 78–83 (2020). [doi:10.1038/s41594-019-0357-0](https://doi.org/10.1038/s41594-019-0357-0) [Medline](#)
  23. J. Pallesen, N. Wang, K. S. Corbett, D. Wrapp, R. N. Kirchdoerfer, H. L. Turner, C. A. Cottrell, M. M. Becker, L. Wang, W. Shi, W.-P. Kong, E. L. Andres, A. N. Kettenbach, M. R. Denison, J. D. Chappell, B. S. Graham, A. B. Ward, J. S. McLellan, Immunogenicity and structures of a rationally designed prefusion MERS-CoV spike antigen. *Proc. Natl. Acad. Sci. U.S.A.* **114**, E7348–E7357 (2017). [doi:10.1073/pnas.1707304114](https://doi.org/10.1073/pnas.1707304114) [Medline](#)
  24. D. Wrapp, N. Wang, K. S. Corbett, J. A. Goldsmith, C.-L. Hsieh, O. Abiona, B. S. Graham, J. S. McLellan, Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *Science* **367**, 1260–1263 (2020). [doi:10.1126/science.abb2507](https://doi.org/10.1126/science.abb2507) [Medline](#)
  25. Y. Lin, Y. Gu, S. A. Wharton, L. Whittaker, V. Gregory, X. Li, S. Metin, N. Cattle, R. S. Daniels, A. J. Hay, J. W. McCauley, Optimisation of a micro-neutralisation assay and its application in antigenic characterisation of influenza viruses. *Influenza Other Respir. Viruses* **9**, 331–340 (2015). [doi:10.1111/irv.12333](https://doi.org/10.1111/irv.12333) [Medline](#)
  26. L. H. Evans, R. P. Morrison, F. G. Malik, J. Portis, W. J. Britt, A neutralizable epitope common to the envelope glycoproteins of ecotropic, polytropic, xenotropic, and amphotropic murine leukemia viruses. *J. Virol.* **64**, 6176–6183 (1990). [doi:10.1016/j.vi.64.12.6176-6183](https://doi.org/10.1016/j.vi.64.12.6176-6183) [Medline](#)
  27. M. Hoffmann, H. Kleine-Weber, S. Schroeder, N. Krüger, T. Herrler, S. Erichsen, T. S. Schiergens, G. Herrler, N.-H. Wu, A. Nitsche, M. A. Müller, C. Drosten, S. Pöhlmann, SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. *Cell* **181**, 271–280.e8 (2020). [doi:10.1016/j.cell.2020.02.052](https://doi.org/10.1016/j.cell.2020.02.052) [Medline](#)
  28. X. Ou, Y. Liu, X. Lei, P. Li, D. Mi, L. Ren, L. Guo, R. Guo, T. Chen, J. Hu, Z. Xiang, Z. Mu, X. Chen, J. Chen, K. Hu, Q. Jin, J. Wang, Z. Qian, Characterization of spike glycoprotein of SARS-CoV-2 on virus entry and its immune cross-reactivity with SARS-CoV. *Nat. Commun.* **11**, 1620 (2020). [doi:10.1038/s41467-020-15562-9](https://doi.org/10.1038/s41467-020-15562-9) [Medline](#)
  29. A. C. Walls, Y. J. Park, M. A. Tortorici, A. Wall, A. T. McGuire, D. Velesler, Structure, Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein. *Cell* **181**, 281–292.e6 (2020). [doi:10.1016/j.cell.2020.02.058](https://doi.org/10.1016/j.cell.2020.02.058) [Medline](#)
  30. P. Zhou, X. L. Yang, X. G. Wang, B. Hu, L. Zhang, W. Zhang, H. R. Si, Y. Zhu, B. Li, C. L. Huang, H. D. Chen, J. Chen, Y. Luo, H. Guo, R. D. Jiang, M. Q. Liu, Y. Chen, X. R. Shen, X. Wang, X. S. Zheng, K. Zhao, Q. J. Chen, F. Deng, L. L. Liu, B. Yan, F. X. Zhan, Y. Y. Wang, G. F. Xiao, Z. L. Shi, A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* **579**, 270–273 (2020). [doi:10.1038/s41586-020-2012-7](https://doi.org/10.1038/s41586-020-2012-7) [Medline](#)
  31. K. Wang, W. Chen, Y.-S. Zhou, J.-Q. Lian, Z. Zhang, P. Du, L. Gong, Y. Zhang, H.-Y. Cui, J.-J. Geng, B. Wang, X.-X. Sun, C.-F. Wang, X. Yang, P. Lin, Y.-Q. Deng, D. Wei, X.-M. Yang, Y.-M. Zhu, K. Zhang, Z.-H. Zheng, J.-L. Miao, T. Guo, Y. Shi, J. Zhang, L. Fu, Q.-Y. Wang, H. Bian, P. Zhu, Z.-N. Chen, SARS-CoV-2 invades host cells via a novel route: CD147-spike protein. *bioRxiv* 988345 [Preprint]; 14 March 2020; <https://doi.org/10.1101/2020.03.14.988345>
  32. L. Cantuti-Castelvetri, R. Ojha, L. D. Pedro, M. Djannatian, J. Franz, S. Kuivanen, K. Kallio, T. Kaya, M. Anastasina, T. Smura, L. Levanov, L. Szivovics, A. Tobi, H. Kallio-Kokko, P. Österlund, M. Joensuu, F. A. Meunier, S. Butcher, M. S. Winkler, B. Mollenhauer, A. Helenius, O. Gokce, T. Teesalu, J. Hepojoki, O. Vapalahti, C. Stadelmann, G. Balistreri, M. Simons, Neuropilin-1 facilitates SARS-CoV-2 cell entry and provides a possible pathway into the central nervous system. *bioRxiv* 137802 [Preprint]. 15 July 2020; <https://doi.org/10.1101/2020.06.07.137802>
  33. J. L. Daly, B. Simonetti, C. Antón-Plágaro, M. Kavanagh Williamson, D. K. Shoemark, L. Simón-Gracia, K. Klein, M. Bauer, R. Hollandi, U. F. Greber, P. Horvath, R. B. Sessions, A. Helenius, J. A. Hiscox, T. Teesalu, D. A. Matthews, A. D. Davidson, P. J. Cullen, Y. Yamauchi, Neuropilin-1 is a host factor for SARS-CoV-2 infection. *bioRxiv* 134114 [Preprint]. 5 June 2020; <https://doi.org/10.1101/2020.06.05.134114>
  34. M. C. Freeman, C. T. Peek, M. M. Becker, E. C. Smith, M. R. Denison, Coronaviruses induce entry-independent, continuous macropinocytosis. *mBio* **5**, e01340–e14 (2014). [doi:10.1128/mBio.01340-14](https://doi.org/10.1128/mBio.01340-14) [Medline](#)
  35. T. C. Nash, T. M. Gallagher, M. J. Buchmeier, MHV-1 independent cell-cell spread of mouse hepatitis virus infection requires neutral pH fusion. *Adv. Exp. Med. Biol.* **380**, 351–357 (1995). [doi:10.1007/978-1-4615-1899-0\\_57](https://doi.org/10.1007/978-1-4615-1899-0_57) [Medline](#)
  36. A. Grifoni, J. Sidney, Y. Zhang, R. H. Scheuermann, B. Peters, A. Sette, A Sequence Homology and Bioinformatic Approach Can Predict Candidate Targets for Immune Responses to SARS-CoV-2. *Cell Host Microbe* **27**, 671–680.e2 (2020). [doi:10.1016/j.chom.2020.03.002](https://doi.org/10.1016/j.chom.2020.03.002) [Medline](#)
  37. A. G. Wrobel, D. J. Benton, P. Xu, C. Rouston, S. R. Martin, P. B. Rosenthal, J. J. Skehel, S. J. Gamblin, SARS-CoV-2 and bat RaTG13 spike glycoprotein structures inform on virus evolution and furin-cleavage effects. *Nat. Struct. Mol. Biol.* **27**, 763–767 (2020). [doi:10.1038/s41594-020-0468-7](https://doi.org/10.1038/s41594-020-0468-7) [Medline](#)
  38. Y. Cai, J. Zhang, T. Xiao, H. Peng, S. M. Sterling, R. M. Walsh Jr., S. Rawson, S. Rits-Volloch, B. Chen, Distinct conformational states of SARS-CoV-2 spike protein. *Science* **369**, 1586–1592 (2020). [doi:10.1126/science.abb4251](https://doi.org/10.1126/science.abb4251) [Medline](#)
  39. B. Turoňová, M. Sikora, C. Schürmann, W. J. H. Hagen, S. Welsch, F. E. C. Blanc, S. von Bülow, M. Gecht, K. Bagola, C. Hörner, G. van Zandbergen, J. Landry, N. T. D. de Azevedo, S. Mosalaganti, A. Schwarz, R. Covino, M. D. Mühlebach, G. Hummer, J. Krijnse Locker, M. Beck, In situ structural analysis of SARS-CoV-2 spike reveals flexibility mediated by three hinges. *Science* **370**, 203–208 (2020). [doi:10.1126/science.abb5223](https://doi.org/10.1126/science.abb5223) [Medline](#)
  40. S. Klein, M. Cortese, S. L. Winter, M. Wachsmuth-Melm, C. J. Neufeldt, B. Cerikan, M. L. Stanifer, S. Boulant, R. Bartenschlager, P. Chlanda, SARS-CoV-2 structure and replication characterized by in situ cryoelectron tomography. *bioRxiv* 167064 [Preprint]. 16 August 2020; <https://doi.org/10.1101/2020.06.23.167064>

#### ACKNOWLEDGMENTS

We thank L. James and J. Luptak for the SARV CoV2 N expression construct and M. Pizzato for the SARS CoV2 S cDNA. We also thank the entire CRICK COVID-19 Consortium. We are grateful for assistance from the Cell Services and High Throughput Screening facilities at the Francis Crick Institute and UCLH Biochemistry (A. Goyale and C. Wilson) and to Mr Michael Bennet and Mr Simon Caidan for training and support in the high-containment laboratory. **Funding:**

This work was supported by a Centre of Excellence Centre for Adolescent Rheumatology Versus Arthritis grant, 21593, as well as support from the Great Ormond Street Childrens Charity, CureJM Foundation and the NIHR Biomedical Research Centres at GOSH and UCLH. This work was supported by the Francis Crick Institute, which receives its core funding from Cancer Research UK, the UK Medical Research Council, and the Wellcome Trust. **Author contributions:** Experimental design, C.C., L.R.W., R.B., C.S., S.G., B.S., J.McC., S.J.G., L.E.McC., P.C., E.N., and G.K. Investigation, K.W.N., N.F., G.H.C., A.Ro., R.H., S.H., R.U., C.E., A.G.W., D.J.B., C.R., W.B., R.T., A.A.-D., P.H., and D.J. Reagents and Samples, J.H., H.R., S.P., C.F.H., K.T., E.S., G.Y.S., M.J.S., P.A.W., C.M., B.R.J., M.G.Li.W., L.R.M., E.C.R., A.Ra., and H.P. Writing, L.E.McC., P.C., E.N., and G.K., with contributions from C.C., L.R.W., K.W.N., N.F., and G.H.C. Supervision, N.O'R., S.K., A.Ri., C.C., L.R.W., R.B., C.S., S.G., B.S., J.McC., S.J.G., L.E.McC., P.C., E.N., and G.K. **Competing interests:** The authors declare no competing interests. **Data and materials availability:** All data are available in the main text or the supplementary materials. This work is licensed under a Creative Commons Attribution 4.0 International (CC BY 4.0) license, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. To view a copy of this license, visit <https://creativecommons.org/licenses/by/4.0/>. This license does not apply to figures/photos/artwork or other content included in the article that is credited to a third party; obtain authorization from the rights holder before using such material.

## SUPPLEMENTARY MATERIALS

[science.sciencemag.org/cgi/content/full/science.abe1107/DC1](http://science.sciencemag.org/cgi/content/full/science.abe1107/DC1)

Materials and Methods

Supplementary Text

Figs. S1 to S18

Tables S1 and S2


References (18–40)

MDAR Reproducibility Checklist


31 July 2020; accepted 29 October 2020

Published online 6 November 2020


10.1126/science.abe1107



**Fig. 1. Flow cytometric detection and specificity of antibodies reactive with SARS-CoV-2 S.** (A) Detection of IgG, IgA, and IgM in five individuals from each indicated group. IgM levels are indicated by a heatmap. (B to D) Inhibition of SARS-CoV-2 S binding of sera from SARS-CoV-2-infected (SARS-CoV-2<sup>+</sup>, n=10) or SARS-CoV-2-uninfected (SARS-CoV-2<sup>-</sup> HCoV<sup>+</sup>, n=6) patients by soluble S1 or S2. Flow cytometry profile of one representative patient per group is shown (B). Mean frequency of positive cells (C). \**P*=0.015; \*\**P*=0.006, one-way analysis of variance (ANOVA) on Ranks. Mean staining intensity (mean fluorescent intensity (MFI) of sample as a percentage of negative control MFI) (D). In C and D, dots represent individual samples from one of three similar experiments.




**Fig. 2. Prevalence of SARS-CoV-2 S-cross-reactive antibodies detected by different methods.** (A) Flow cytometry and ELISA results for each sample in cohorts A and C to E (table S1). (B) Flow cytometry and ELISA results for serum samples from SARS-CoV-2-uninfected pregnant women. (C to E) SARS-CoV-2 S-cross-reactive antibodies in healthy children and adolescents. Representative flow cytometry profiles of seronegative donors (Negative) or COVID-19 patients (Positive) and of SARS-CoV-2-uninfected adolescents with SARS-CoV-2-cross-reactive antibodies (C). Frequency of cells stained with all three antibody classes (IgG<sup>+</sup>IgM<sup>+</sup>IgA<sup>+</sup>) or only with IgG (IgG<sup>+</sup>), ranked by their IgG<sup>+</sup>IgM<sup>+</sup>IgA<sup>+</sup> frequency (D). The dashed line denotes the assay sensitivity cut-off. Flow cytometry and ELISA results for each sample (E). (F) Prevalence of SARS-CoV-2 S-cross-reactive antibodies in the indicated age groups (line) and frequency of cells that stained only with IgG (dots) in all samples for which the date of birth was known. The heatmaps in A, B, and E represent the quartile values above each assay's technical cut-off.



**Fig. 3. Neutralization of SARS-CoV-2 S pseudotypes and authentic SARS-CoV-2 by SARS-CoV-2-infected and -uninfected patient sera. (A)** Inhibition of transduction efficiency of SARS-CoV-2 S or VSVg pseudotypes by adult COVID-19 patients who seroconverted (SARS-CoV-2<sup>+</sup> Adults Ab<sup>+</sup>) or not (SARS-CoV-2<sup>+</sup> Adults Ab<sup>-</sup>), and SARS-CoV-2-uninfected adult (SARS-CoV-2<sup>-</sup> Adults Ab<sup>+</sup>) or children and adolescent donors (SARS-CoV-2<sup>-</sup> Children/Adolescents Ab<sup>+</sup>) donors with SARS-CoV-2 S-binding antibodies. Each line is an individual serum sample. **(B)** Authentic SARS-CoV-2 neutralization titers of sera from the same donors as in A, as well as SARS-CoV-2-uninfected donors without SARS-CoV-2 S-binding antibodies (Ab<sup>-</sup>). Dots represent individual samples. \* $P=0.037$ ; \*\* $P=0.014$ ; ns: not significant, one-way ANOVA on Ranks.





**Fig. 4. Mapping of cross-reactive epitopes in SARS-CoV-2 S.** (A) Signal intensity for each overlapping peptide along the length of SARS-CoV-2 S covered in the peptide arrays, using pooled sera with (Ab<sup>+</sup>) or without (Ab<sup>-</sup>) flow cytometry-detectable SARS-CoV-2 S-reactive antibodies. Differentially recognized peaks are boxed. (B) Alignment of the amino acid sequences of SARS-CoV-2 and HCoV S glycoproteins. Boxes indicate predicted core epitopes. (C) Mapping of predicted epitopes targeted on the trimeric SARS-CoV-2 spike. The S1 (blue) and S2 (pink) of one monomer are colored. Epitopes are shown for one monomer; the circled dashed line represent the membrane proximal region not present in the structure. (D) (Left), Reactivity with the S glycoproteins of each HCoV of the indicated sera with (Ab<sup>+</sup>) or without (Ab<sup>-</sup>) flow cytometry-detectable SARS-CoV-2 S-reactive antibodies, determined by flow cytometry. Each column is an individual sample. Rows depict the staining for each antibody class. (Right), Correlation coefficients between percentages of IgG staining for SARS-CoV-2 S and IgG, IgM, and IgA staining for each HCoV S glycoprotein.

## Preexisting and de novo humoral immunity to SARS-CoV-2 in humans

Kevin W. Ng, Nikhil Faulkner, Georgina H. Cornish, Annachiara Rosa, Ruth Harvey, Saira Hussain, Rachel Ulferts, Christopher Earl, Antoni G. Wrobel, Donald J. Benton, Chloe Roustán, William Bolland, Rachael Thompson, Ana Agua-Doce, Philip Hobson, Judith Heaney, Hannah Rickman, Stavroula Paraskevopoulou, Catherine F. Houlihan, Kirsty Thomson, Emilie Sanchez, Gee Yen Shin, Moira J. Spyer, Dhira Joshi, Nicola O'Reilly, Philip A. Walker, Svend Kjaer, Andrew Riddell, Catherine Moore, Bethany R. Jebson, Meredyth Wilkinson, Lucy R. Marshall, Elizabeth C. Rosser, Anna Radziszewska, Hannah Peckham, Coziana Ciurtin, Lucy R. Wedderburn, Rupert Beale, Charles Swanton, Sonia Gandhi, Brigitta Stockinger, John McCauley, Steve J. Gamblin, Laura E. McCoy, Peter Cherepanov, Eleni Nastouli and George Kassiotis

published online November 6, 2020

### ARTICLE TOOLS

<http://science.sciencemag.org/content/early/2020/11/05/science.abe1107>

### SUPPLEMENTARY MATERIALS

<http://science.sciencemag.org/content/suppl/2020/11/05/science.abe1107.DC1>

### RELATED CONTENT

<http://stm.sciencemag.org/content/scitransmed/12/564/eabd5487.full>  
<http://stm.sciencemag.org/content/scitransmed/12/550/eabc3539.full>  
<http://stm.sciencemag.org/content/scitransmed/12/555/eabc9396.full>  
<http://stm.sciencemag.org/content/scitransmed/12/557/eabc5332.full>

### REFERENCES

This article cites 39 articles, 17 of which you can access for free  
<http://science.sciencemag.org/content/early/2020/11/05/science.abe1107#BIBL>

### PERMISSIONS

<http://www.sciencemag.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of Service](#)

*Science* (print ISSN 0036-8075; online ISSN 1095-9203) is published by the American Association for the Advancement of Science, 1200 New York Avenue NW, Washington, DC 20005. The title *Science* is a registered trademark of AAAS.

Copyright © 2020 The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works. Distributed under a Creative Commons Attribution License 4.0 (CC BY).